

**New Nucleotide Sequences Coding for the thrE Gene and
Process for the Enzymatic Production of L-threonine using
Coryneform Bacteria**

Patent Claims

- 5 1. Preferably recombinant DNA derived from Corynebacterium
and replicable in coryneform microorganisms, which
contains at least one nucleotide sequence that codes
for the thrE gene.
2. Replicable DNA according to claim 1 with
- 10 (i) the nucleotide sequences shown in SEQ-ID-No. 1,
or SEQ-ID No. 3, which code for the thrE gene,
or
- (ii) at least once sequence that corresponds to the
sequences (i) within the degeneration region of
- 15 (iii) at least once sequence that hybridises with the
sequences complementary to the sequences (i) or
(ii), and/or optionally
- (iv) functionally neutral sense mutations in (i).
- 20 3. Amino acid sequence of the protein, derived from the
nucleotide sequences according to claim 1 or 2, shown
in SEQ-ID-No. 2 and in SEQ-ID-No. 4.
4. Coryneforme microorganisms, in particular of the genus
Corynebacterium, transformed by the introduction of one
- 25 or more of the replicable DNA according to claim 1 or
2.
5. Corynebacterium glutamicum DM368-2 pZ1thrE, filed under
Number DSM 12840.
6. Process for producing L-threonine by fermentation of
- 30 coryneform bacteria, characterised in that bacteria are
used in which nucleotide sequences coding for the thrE

096351-09304
T0220-12569660

gene are amplified, and in particular are overexpressed.

7. Process according to claim 6, characterised in that bacteria are used in which in addition one or more genes of the threonine biosynthesis pathway is/are amplified.
8. Process according to claims 6 and 7, characterised in that a strain transformed with a plasmid vector is used and the plasmid vector carries the nucleotide sequence coding for the thrE gene.
9. Process according to claims 6 and 8, characterised in that the thrE gene is overexpressed in microorganisms that contain further metabolite or antimetabolite resistance mutations.
10. Process according to claims 6 to 9, characterised in that the microorganisms in order to achieve over-expression are fermented in altered culture media, and/or the fermentation conditions are changed.
11. Process according to claims 6 to 10, characterised in that microorganisms are used in which the metabolic pathways that reduce threonine formation are at least partially switched off.
12. Process according to claims 6 to 11, characterised in that microorganisms are used in which in addition to the thrE gene the remaining genes of the metabolic pathway for threonine formation are amplified individually or jointly (overexpressed).
13. Process for producing L-threonine, characterised in that the following steps are carried out:
- a) Fermentation of microorganisms according to one or more of the preceding claims, in which at least the thrE gene is amplified (overexpressed) optionally in combination with further genes,

- 90 | 9 ||||| 00 | 00

10

15

b) in the case of transposon mutagenesis is selected for the transposon preferably exhibiting resistance to antibiotics, and the thrE gene is thereby obtained.